



Recombinant Human Vascular Endothelial Growth Factor₁₆₅
(rhVEGF₁₆₅)
Catalog Number: 105-05

Description	VEGF is a potent growth and angiogenic cytokine. It stimulates proliferation and survival of endothelial cells, and promotes angiogenesis and vascular permeability. Expressed in vascularized tissues, VEGF plays a prominent role in normal and pathological angiogenesis. Substantial evidence implicates VEGF in the induction of tumor metastasis and intra-ocular neovascular syndromes. VEGF signals through the three receptors; fms-like tyrosine kinase (flt-1), KDR gene product (the murine homolog of KDR is the flk-1 gene product) and the flt4 gene product.
Synonyms	
AA Sequence	APMAEGGGQN HHEVVKFMDV YQRSYCHPIE TLVDIFQEYP DEIEYIFKPS CVPLMRCGGC CNDEGLECVP TEESNITMQI MRIKPHGGQH IGEMSFLQHN KCECRPKKDR ARQENPCGPC SERRKHLFVQ DPQTCKCCK NTDSRCKARQ LELNERTCRC DKPRR
Source	<i>Escherichia coli</i>
Molecular Weight	Approximately 38.2 kDa disulfide-linked homodimeric protein consisting of two 165 amino acid polypeptide chains.
Purity	>95% by SDS-PAGE and HPLC analyses.
Biological Activity	Fully biologically active. The ED ₅₀ is 1-8ng/ml, as determined by HUVEC cell proliferation.
Physical Appearance	White lyophilized powder.
Formulation	Lyophilized from a 0.2µm filtered solution in PBS, pH 7.4.
Endotoxin	< 1EU/µg of growth factor as determined by LAL method.
Reconstitution	Reconstitute in sterile distilled water containing 0.1% BSA to a concentration of 0.1-1.0 mg/mL.
Storage	Store at -20°C after receiving. Upon reconstitution, store at 2-8°C for up to one week. For maximal stability, aliquot and store at -20°C. Avoid repeated freeze/ thaw cycles.
Usage	This product is for research use only. It is not approved for use in humans, animals, or <i>in vitro</i> diagnostic procedures.