

Human Neuron-hippocampal (HN-h) Lysate Catalog Number: 1546(200 µg)/1546-4(400 µg)

Product Information

Quantity:	200 µg/400 µg	
Concentration:	2 mg/ml/4 mg/ml	
Storage buffer:	1X reducing sample buffer (50 mM Tris-HCl pH 6.8, 5% glycerol, 1% SDS, 0.002% bromphenol blue, 5% beta- mercaptoethanol)	250k Da —
Lysis buffer:	Modified RIPA buffer (50 mM Tris-HCl pH 7.2, 150 mM NaCl, 1% NP-40, 1mM EDTA, 1 mM EGTA, 0.4 mM PMSF, 5 µg/ml aprotinin, 5 µg/ml leupeptin, 1 µg/ml pepstatin, 1 mM Na ₃ VO ₄ , 5 mM NaF)	130k Da — 100k Da — 70k Da —
Quality control:	Lysate quality was assured with a NuPAGE 4-12% Bis-Tris Gel stained with Coomassie brilliant blue.	55k Da —

Storage/Handling

Stable for 6 months at -20°C from date of shipment. For maximum recovery of product, centrifuge the original vial after thawing and before removing the cap. Aliquot to avoid repeated freezing and thawing.

Application

Cell lysate need to to be boiled for 5 minutes before loading on SDS-PAGE for western blot. 40 μ g per lane is recommended for mini gel.

QC SDS-PAGE gel

35k Da

27k Da

15k Da 10k Da